

# GEOGRAPHIC DISTRIBUTION

## SERPENTES

**GRAYIA THOLLONI** (Thollon's Water Snake). SENEGAL: CAP-VERT REGION: Sebikotane, 14°45'N, 17°08'W. 1963. R. P. Berhaut. Verified by Ivan Ineich. Museum National d'Histoire Naturelle at Paris (MNHN 1964. 145). Second record for Senegal (Trape and Mane 1995, Herpetol. Rev. 26:156) and third record for West Africa W of Nigeria (Hughes 1983, Bonn. zool. Beitr. 34:311–356; Dunger 1972, Nigerian Field 37(1):21–38). Specimen was previously erroneously determined as *Grayia smithii* on the MNHN register by Villiers who gave “North Nigeria to Lake Tanganyika” as distribution for the species (Villiers 1975, Les Serpents de l'Ouest Africain. Institut Fondamental d'Afrique Noire, Init. Etudes Afr. n°2:1–195, 3rd ed.). This new locality is situated 130 km (airline) NW of the previous northwesternmost known locality.

Submitted by **OLIVER PAUWELS**, Museum National d'Histoire Naturelle, Laboratoire de Zoologie (Reptiles et Amphibiens), 25 rue Cuvier, F-75005 Paris, France, or Koninklijk Museum voor Midden-Africa, Vertebrate Section (Herpetology), Steenweg op Leuven, B-3080 Tervuren, Belgium.